

FITC-Linked Polyclonal Antibody to Immunoglobulin G1 (IgG1)

Catalog No: FY-AB33587

Organism Species: Homo sapiens (Human)

Application: WB; IHC; ICC; IF.

Alternative Names: IGHG1; Ig Gamma-1 Chain C Region; Immunoglobulin Heavy Constant Gamma 1; G1m Marker; Immunoglobulin Gm1

PROPERTIES

Source	Antibody labeling
Host species	293F
Cross Reactivity	-
Purification	Antigen-specific + Protein A affinity chromatography
Research Area	Immune molecule;
Appearance	Liquid
Size	200μl;500μg/mL
Formulation	0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.
Immunogen	Polyclonal Antibody to Immunoglobulin G1 (IgG1)
Application	Western blotting: 0.5-2μg/mL
	Immunocytochemistry in formalin fixed cells: 5-20μg/mL
	Immunohistochemistry in formalin fixed frozen section: 5-20μg/mL
Storage instructions	Stable for 12 months. at -20°C from date of shipment. Aliquot to avoid repeated freezing and thawing. Store at 2-8°C for frequent use. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap.
Stability Test	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

NOTE:The product listed herein is for research use only and is not intended for use in human or clinical diagnosis. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license. We cannot be responsible for patent infringements or other violations that may occur with the use of this product. The product listed herein is for research use only and is not intended for use in human or clinical diagnosis. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license. We cannot be responsible for patent infringements or other violations that may occur with the use of this product.