

Anti-Fc Fragment Of IgG Low Affinity IIIb Receptor (FcgR3B) Polyclonal Antibody

Catalog No: FY-AB54001

Organism Species: Homo sapiens (Human)

Application: Flow cytometry.

Alternative Names: CD16b; CD16-B; FCGR3-B; FCG3; FcgRIII; Low affinity immunoglobulin gamma Fc region receptor III-

В

PROPERTIES

-
Rabbit
-
Antigen-specific + Protein A affinity chromatography
CD & Adhesion molecule;
Liquid
50test;500μg/mL
0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.
-
For flow cytometry, the suggested use of this reagent is 1-5 μ l per 10^6 cells in 100 μ l volume.
Optimal working dilutions could be determined by end user.
Stable for 12 months. at -20°C from date of shipment. Aliquot to avoid repeated freezing and thawing.
Store at 2-8°C for frequent use. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap.
The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

NOTE: The product listed herein is for research use only and is not intended for use in human or clinical diagnosis. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license. We cannot be responsible for patent infringements or other violations that may occur with the use of this product. The product listed herein is for research use only and is not intended for use in human or clinical diagnosis. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license. We cannot be responsible for patent infringements or other violations that may occur with the use of this product.

www.feiyuebio.com Tel: (0086)027-87002654 Fax: (0086)027-878002654