

FITC-Linked Anti-Tumor Necrosis Factor Receptor Superfamily, Member 7 (TNFRSF7) Monoclonal Antibody

Catalog No: FY-AB54079

Organism Species: Homo sapiens (Human)

Application: Flow cytometry.

Alternative Names: CD27; S152; T14; Tp55; CD27L receptor;

PROPERTIES

Source	-
Host species	293F
Cross Reactivity	-
Purification	Protein A+G
Research Area	Signal transduction;CD & Adhesion molecule;Apoptosis;Immune molecule;
Appearance	Liquid
Size	50test;1mg/mL
Formulation	0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.
Immunogen	-
Application	For flow cytometry, the suggested use of this reagent is 1-5 μ l per 10^6 cells in 100 μ l volume.
	Optimal working dilutions could be determined by end user.
Storage instructions	Stable for 12 months. at -20°C from date of shipment. Aliquot to avoid repeated freezing and thawing. Store at 2-8°C for frequent use. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap.
Stability Test	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

NOTE:The product listed herein is for research use only and is not intended for use in human or clinical diagnosis. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license. We cannot be responsible for patent infringements or other violations that may occur with the use of this product. The product listed herein is for research use only and is not intended for use in human or clinical diagnosis. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license. We cannot be responsible for patent infringements or other violations that may occur with the use of this product.